arXiv:1305.0139v1 [math.PR] 1 May 2013

Condensation of a two-dimensional random walk and the Wulff crystal

by Nathanaël Berestycki* Ariel Yadin†

May 2, 2013

Abstract

We introduce a Gibbs measure on nearest-neighbour paths of length t in the Euclidean planar lattice, where each path is penalized by a factor proportional to the size of its boundary and an inverse temperature β. This measure gives a random walk description of the Wulff crystal, representing the distribution of a diluted polymer in a poor solvent. We prove that in two dimensions, if the inverse temperature β is larger than the logarithm of the connective constant, then the random walk condensates to a set of diameter $t^{1/3}$, up to logarithmic corrections. We further speculate that the limiting shape shares the same exponents as the KPZ universality class. A similar result holds for a random walk conditioned to have local time greater than β everywhere in its range.

Keywords: Gibbs measure, condensation, self–attractive random walk, Wulff crystal, large deviations, Donsker–Varadhan principle.

MSC 2010 classification: 60K35, 60J27, 60F10

---

*University of Cambridge. Supported in part by EPSRC grants EP/GO55068/1 and EP/I03372X/1
†Ben Gurion University of the Negev.



Figure 1: Simulations using a Gibbs sampler algorithm of a random walk cluster with $t = 25{,}000$ steps, corresponding to four different values of β: $\beta = 0.01$, $\beta = 0.1$, $\beta = 1$, and $\beta = 2$. Pictures realised by Tom Begley.

Contents

1 Introduction **2**
- 1.1 Statement of the main results 2
- 1.2 Variants 4
- 1.3 Notation 5

2 Change of measure estimates **6**
- 2.1 Statement of general theorem 6
- 2.2 Canonical paths 7
- 2.3 Flows and hitting estimates 9
- 2.4 Concentration estimates for local time 12
- 2.5 Change of measure 15

3 Proof of Theorem 1.1 **17**
- 3.1 Partition function estimate 17
- 3.2 Discrete planar isoperimetry 20
- 3.3 Proof of condensation 20
- 3.4 Proof of Theorems 1.3 and 1.4 23

4 Open problems and conjectures **24**

1 Introduction

1.1 Statement of the main results

Let $d \geq 1$ and let Ω be the space of nearest-neighbours right-continuous infinite paths $(\omega_t, t \in [0, \infty))$ on $\mathbb{Z}^d$ started at $\omega_0 = 0$, and let $(X_t(\omega), t \geq 0)$ be the canonical process. Let $\mathbb{P}$ denote the law of simple random walk on $(\Omega, \mathcal{F})$ in continuous time with rate one, where $\mathcal{F}$ denotes the σ-field generated by X.

Our main result deals with geometric properties of some penalizations of random walks on $\mathbb{Z}^d$ by their boundary. More precisely, we introduce a Gibbs measure μ on random paths defined as follows. Let $R_t = \{v \in \mathbb{Z}^d : X_s = v \text{ for some } s \leq t\}$ denote the *range* of the walk at time t. For a given time t, we consider the Hamiltonian H given by

$$H(\omega) = |\partial R_t|, \tag{1.1}$$

where for a set G, $\partial G = \{x \in G : x \sim y \text{ for some } y \notin G\}$ is the (inner) vertex-boundary of G. The associated Gibbs measure on random paths $\mu = \mu_t$ is obtained by considering the measure μ defined by

$$\frac{d\mu}{d\mathbb{P}}(\omega) = \frac{1}{Z}\exp(-\beta H(\omega)) \tag{1.2}$$

on $\mathcal{F}$. Here $\beta > 0$ is a positive number playing the role of inverse temperature and $Z = Z(\beta, t) = \mathbb{E}(\exp(-\beta|\partial R_t|))$ is a normalizing factor called the partition function. In plain words, the Gibbs measure μ penalizes every site on the boundary of the range R_t by a fixed amount $e^{-\beta}$. Hence μ favours "highly condensed" configurations. Interpreting the random walk $(X_0, \dots, X_t)$ as a chain of t monomers, the Gibbs measure μ represents the law of a diluted polymer in a poor solvent. With a constraint on the maximal volume and a boundary penalization, the random set R_t may be seen as a random walk construction of the Wulff crystal droplet with inverse temperature β. The Wulff construction ([24]) is a method to determine the equilibrium shape of crystals based on surface energy minimization. In the percolation context, rigorous derivation of the limiting shape were first given by Alexander, Chayes and Chayes [2] in two dimensions, while in the context of the Ising model, this was achieved in a nearly simultaneous and independent way in a celebrated work of Dobrushin, Kotecký and Shlosman [8]. See [4] for a recent monograph giving a detailed overview of the subject.

We will be interested in describing the geometry of the shape of R_t in the low temperature regime, where β is sufficiently large. Our main result gives precise estimates for the condensation effect that results from the self-interaction in dimension 2. If $G \subset \mathbb{Z}^2$, let $\operatorname{diam} G$ denote the (Euclidean) diameter of G:

$$\operatorname{diam} G = \sup\{|z - w|, z, w \in G\}.$$

Theorem 1.1. *Assume $d = 2$. Let α be the connective constant of $\mathbb{Z}^2$ and let $\beta_0 = \log \alpha$. Then for all $\beta > \beta_0$,*

$$\mu\left\{\frac{1}{\ell_1(t)}t^{1/3} \leq \operatorname{diam}(R_t) \leq \ell_2(t)t^{1/3}\right\} \to 1, \tag{1.3}$$

as $t \to \infty$, where $\ell_1(t) = c_1(\log t)^{5/2}$ and $\ell_2(t) = c_2(\log t)^5$ and the constants c_1, c_2 depend on β only.

Remark 1.2. *The exponent $1/3$ in Theorem 1.1 is in agreement with the physics predictions, which asserts that the radius of gyration has exponent $\nu = 1/3$.*

The proof gives precise estimates on the probability of these events, as well as estimates on the partition function $Z = Z(\beta, t)$. We refer the reader to Theorem 3.6 for a precise statement, and refer to Open Problem 3 and the following paragraph for a precise conjecture on the geometry of R_t when $d = 2$.

A related problem was studied by E. Bolthausen [3]. In that work, the energy $H(\omega)$ serving to define the Gibbs measure μ in (1.2) is taken to be $\hat{H}(\omega) = |R_t(\omega)|$, the size of the range (as opposed to that of its boundary). Thus $d\hat{\mu} = \hat{Z}_t^{-1} \exp(-\beta \hat{H}(\omega)) d\,\mathbb{P}$. Bolthausen's result is that the random walk condensates to a set of diameter $t^{1/4}$, which is close in the Hausdorff sense to a Euclidean ball of that diameter. In both problems, good bounds on the partition functions Z_t and $\hat{Z}_t$ play a crucial role. In the case where the energy is just the volume, we have $\hat{Z}_t = \mathbb{E}(\exp(-\beta|R_t|))$, and precise asymptotics for this quantity were already obtained by Donsker and Varadhan [13]. This is a considerably easier problem than the one considered here, essentially because $|R_t(\omega)|$ is "almost" a continuous function of its local time profile, viewed as a probability measure on $\mathbb{Z}^d$. In particular, the powerful machinery of large deviations theory provides the right tools to study this question. This goes a long way in explaining the appearance of the Euclidean ball as a limit shape, and explains why the inverse temperature β is not a relevant parameter in that model.

In contrast, here we believe that the limit shape depends on β and is not a rotationally symmetric ball, and may even contain some flat faces. So the microscopic geometry of the lattice is important even to determine the macroscopic shape of the random walk cluster, and thus there is no hope in directly applying the large deviations machinery to the problem. Instead, a direct and delicate quantitative control of the Markov chain obtained by performing the right change of measure is needed. This is the main technical contribution of this paper.

1.2 Variants

Our technique is sufficiently robust that it yields similar results for a number of models which turn out to be quite closely related.

A first variant consists in taking a slightly different Hamiltonian $\tilde{H}$, defined by

$$\tilde{H}(\omega) = \sum_{x \in \partial R_t} L_t(x), \tag{1.4}$$

Thus the penalisation takes into account not only the size of the boundary, but also the amount of time spent on it. Define $d\tilde{\mu} = (\tilde{Z})^{-1} \exp(-\beta \tilde{H}) d\,\mathbb{P}$ on $\mathcal{F}$.

Theorem 1.3. *Theorem 1.1 still holds true with $\tilde{\mu}$ instead on μ.*

A second variant which we consider looks quite different superficially. Define the event

$$\mathcal{E}_t = \{L_t(x) \geq \beta, \forall\, x \in R_t\}. \tag{1.5}$$

This gives a lower-bound on the density of local time uniformly over the range R_t. This also favours highly condensed configurations. As we will see later on (see for instance (3.14)), the conditioning also heavily penalizes shapes R_t with large boundaries: essentially, every point on the boundary penalizes the shape by a factor of order $e^{-\beta}$, and so we expect a similar behaviour to Theorem 1.1. Indeed, we obtain the following results:

Theorem 1.4. *Assume $d = 2$ and $\beta > \beta_1$ for some $\beta_1 < \infty$. Then*

$$\mathbb{P}\left(\frac{1}{\ell_1(t)} t^{1/3} \leq \operatorname{diam}(R_t) \leq \ell_2(t) t^{1/3} \,\middle|\, \mathcal{E}_t \right) \to 1 \tag{1.6}$$

as $t \to \infty$.

Note that by the isoperimetric inequality, the volume $|R_t|$ is at most $t^{2/3}\ell_2(t)^2$, so the typical local time at a point will be of order $t^{1/3}$ under $\mathbb{P}(\cdot|\mathcal{E}_t)$. This is quite surprising, since in the absence of any conditioning, the local time at a typical vertex is of order $\log t$. Thus requiring a (finite) uniform lower bound β on the local time of each vertex of R_t makes the actual value of the local time at a typical vertex jump from $\log t$ to $t^{1/3}$.

Acknowledgements. This work started when Ariel Yadin was a Herschel Smith postdoctoral fellow at the Statistical Laboratory, University of Cambridge. We gratefully acknowledge the financial support of the Herschel Smith fund and EPSRC grant EP/GO55068/1, without which this work would not have been possible. We thank Itai Benjamini for raising the question of the random walk in Theorem 1.4, which eventually led us to this work. Part of this work was carried out while the first author was visiting Microsoft Research. He is grateful to Yuval Peres for his invitation and interesting related discussions. We also thank Ofer Zeitouni for some very useful conversations regarding large deviations and the link between our work and [6]. We thank Tom Begley for allowing us to use the pictures in Figure 1.

1.3 Notation

Let $d \geq 1$ and let Ω be the space of nearest-neighbours right-continuous infinite paths $(\omega_t, t \in [0,\infty))$ on $\mathbb{Z}^d$ and let $(X_t(\omega), t \geq 0)$ be the canonical process. Let $\mathbb{P}$ denote the law of simple random walk on $(\Omega, \mathcal{F})$ in continuous time with rate one, where $\mathcal{F}$ denotes the σ-field generated by X.

A *conductance* or *weight function* is a non-negative function c on the directed edges of $\mathbb{Z}^d$, that is symmetric (so actually a function of the un-directed edges), such that the total weight at any vertex is finite:

$$c_x = \sum_{y\sim x} c(x,y) < \infty.$$

The canonical continuous time Markov process is that with transition rates given by $c(x,y)$. $\mathbb{P}^c_x, \mathbb{E}^c_x$ denote the probability measure and expectation of this process with respect to a weight function c and started at $X_0 = x$. When omitting the subscript, we refer to $x = 0$. The weight function may also be omitted when it is clear from the context.

If $e = (x,y)$ is an (oriented) edge of $\mathbb{Z}^d$ and ϕ a (real-valued) function on its vertices, denote $\nabla\phi(e) = \nabla\phi(x,y) = \phi(y) - \phi(x)$ the discrete gradient, and

$$\mathcal{E}(\phi,\psi) = \sum_{x\sim y} \nabla\phi(x,y)\nabla\phi(x,y). \tag{1.7}$$

$\mathcal{E}(\phi,\phi)$ is the Dirichlet energy of the function ϕ. We also use the Dirichlet energy in a set S:

$$\mathcal{E}_S(\phi,\psi) = \sum_{\substack{x\sim y\\ x,y\in S}} \nabla\phi(x,y)\nabla\psi(x,y).$$

Let also

$$\mathcal{L}f(x) = \sum_{y\sim x} \nabla f(x,y)$$

denote the discrete Laplacian on $\mathbb{Z}^d$. This is a self adjoint operator with respect to the inner product

$$\langle\phi,\psi\rangle = \sum_x \phi(x)\psi(x).$$

That is, $\langle\mathcal{L}\phi,\psi\rangle = \langle\phi,\mathcal{L}\psi\rangle$.

2 Change of measure estimates

2.1 Statement of general theorem

We start by outlining our strategy. A key difficulty in Theorem 1.1 is to obtain good bounds, and in particular lower bounds, on the partition function $Z_t = \mathbb{E}(\exp(-\beta|\partial R_t|))$. Essentially, the main contribution to this expectation comes from configurations where the typical diameter is of order $t^{1/3}$, and the range has a boundary size which is not much larger (a log factor away). To get a lower bound on the partition function, it suffices to estimate the probability of such events. Thus we will fix a square S of size approximately $t^{1/3}$, and we will estimate the probability that the random walk stays in S and that all but a handful of vertices are visited. The main difficulty arises from the need to ensure that all vertices are indeed visited. This is delicate, especially near the boundary, since each time the walk visits the boundary it shouldn't leave the square. Hence we require that sites near the boundary are visited, but typically only a finite number of times. In order to achieve this, we carefully choose a profile $\pi(x)$, where $\pi(x) > 0$ only in the square S and $\sum_S \pi(x) = 1$, and require that the random walk has a local time profile approximately given by π. We then prove precise quantitative estimates concerning the deviation of the walk from this profile.

We now introduce our framework. Throughout this section we fix a non-negative function $\pi : \mathbb{Z}^d \to \mathbb{R}$ and call $S = \{x \in \mathbb{Z}^d : \pi(x) > 0\}$. We make the following assumptions on π and S: there exists $L > 0$ and some constants $c_i > 0$ independent of L such that:

(B1) π is normalised: $c_1 \le \sum_{x\in S} \pi(x) \le c_2$, and $\max_x \pi(x) \le c_3/L^d$

(B2) For every $z \in S$, there exists a cone $C_z \subset S$ such that z is the apex of C_z, $\operatorname{diam}(C_z) \ge c_4 L$ for some constant c_4 independent of z, and the angle of C_z is uniformly bounded below by some $\alpha > 0$. For all $z \in S$ and all $w \in C_z$,

$$\pi(w) \ge c_5 \pi(z). \tag{2.1}$$

(B3) There exists for each pair of points $x, y \in S$ a fixed path $\gamma_{xy} \subset S$ so that $|\gamma_{xy}| \le L$ and for every edge $e = (e^-, e^+) \in \gamma_{xy}$,

$$\pi(e^\pm) \ge c_6(\pi(x) \wedge \pi(y)). \tag{2.2}$$

Furthermore, for each edge e,

$$\#\{x, y \in S \ : \ e \in \gamma_{xy}\} \le c_7 L^{d+1}. \tag{2.3}$$

(B4) There exists $c_8 > 0$ and c_9 such that $\min_x \pi(x) \ge c_8/L^{c_9}$.

(B5) There exists c_{10} such that $\pi(x)/\pi(y) \le c_{10}$ if $x \sim y$.

Note that since $|\gamma_{xy}| \le L$ for every pair $x, y \in S$, we have $L \ge \operatorname{diam}(S)$. In practice, we will often choose $L = \operatorname{diam}(S)$. Under these assumptions we will prove the following result. Let $f(z) = \sqrt{\pi(z)}$, and define the two following regions:

$$S_- = \{z \in S : \mathcal{L}f(z) < 0\}, S_+ = \{z \in S : \mathcal{L}f(z) > 0\}$$

These are the domains where f is respectively subharmonic or superharmonic. We will prove the following theorem, which is in fact a bit more than needed. Define the event

$$\mathcal{D}_{\alpha,\beta} = \{\alpha\pi(x) \le \frac{L_t(x)}{t} \le \beta\pi(x), x \in \mathbb{Z}^d\}.$$

Let $\mathbb{Q}$ be the Markov chain with transition rates $Q(x,y) = f(y)/f(x)$ if $f(x) > 0$, and $Q(x,y) = 0$ else. Let $Z_t = d\,\mathbb{P}\,/d\mathbb{Q}|_{\mathcal{F}_t}$ denote the Radon-Nikodyn derivative of $\mathbb{P}$ with respect to $\mathbb{Q}$. Then we have the following result.

Theorem 2.1. *Assume (B1)–(B5). Then almost surely on $\mathcal{D}_{\alpha,\beta}$,*

$$\rho \exp(-\frac{\beta}{2}t\mathcal{E}(f,f))) \le Z_t \le \rho^{-1}\exp(-\frac{\alpha}{2}t\mathcal{E}(f,f)). \tag{2.4}$$

Here the error term ρ is given by

$$\rho = \sqrt{\frac{\pi_{\min}}{\pi_{\max}}}\exp(-t(\beta-\alpha)\Phi(f)), \text{ with } \Phi(f) = \sum_{\substack{x\in S_+, y\in S_- \\ x\sim y}} f(x)\nabla f(x,y).$$

In particular, if f is subharmonic, then $\Phi(f) = 0$ and $\rho = \sqrt{\frac{\pi_{\min}}{\pi_{\max}}}$.

Moreover, for all $x, z \in S$, $\mathbb{Q}_x(T_y > t) \le \exp(-ct\pi(y)/(\log L)^3)$, and

$$\mathbb{Q}_x(|L(t,y) - \pi(y)t| \ge \delta\pi(y)t) \le \exp(-c\frac{\delta^2\pi(y)t}{(\log L)^6}). \tag{2.5}$$

Note in particular that

$$\sigma\rho\exp(-\frac{\beta}{2}t\mathcal{E}(f,f))) \le \mathbb{P}_x(D_{\alpha,\beta}) \le \rho^{-1}\exp(-\frac{\alpha}{2}t\mathcal{E}(f,f)) \tag{2.6}$$

where

$$\sigma = 1 - C\sum_{y\in D}\exp(-c\frac{\delta^2\pi(y)t}{(\log L)^6}).$$

where $\delta = \min(|\alpha-1|, |\beta-1|)$. Thus for fixed α, β, we have $\sigma \ge 1/2$ as soon as $t\min_y \pi(y) \ge C(\log L)^7$ for some constant C.

The main idea for the proof of the theorem is to make a change of measure argument so that under the new measure, the random walk has a stationary distribution proportional to φ. Roughly speaking, the role of (B3) is to ensure that the spectral gap under this new measure in D is at least $1/L^2$. On the other hand, the cone condition (B2) ensures that locally the the random walk under this new measure is not "too recurrent", so that it will have at time t a local time profile close reasonably to its limiting value (which is precisely the desired stationary measure, proportional to f).

2.2 Canonical paths

Lemma 2.2. *Let Q be a matrix of rates for a continuous time Markov chain (X_t), on finite state space S with measure $\mathbb{Q}$. Suppose that for some probability vector π we have that $\pi(x)Q(x,y) = \pi(y)Q(y,x)$; that is, π is in detailed balance with Q.*

Let $(\gamma_{xy})_{x,y\in S}$ *be any collection of paths in* S*, one for each pair of vertices. Define*

$$\Theta = \Theta(Q,\pi) = \max_{x\sim y} \frac{1}{\pi(x)Q(x,y)} \sum_{\substack{z,w \\ \gamma_{zw}\ni(x,y)}} \pi(z)\pi(w)|\gamma_{zw}|.$$

Then, for any $\varepsilon > 0$*,*

$$t_{\mathrm{mix}}(\varepsilon) \le -\log\Big(\varepsilon \min_x \pi(x)\Big)\cdot\Theta, \tag{2.7}$$

where

$$t_{\mathrm{mix}}(\varepsilon) = \inf\left\{ t\ge 0 \ : \ \sup_{x,y}\Big|\frac{\mathbb{Q}_x[X_t=y]}{\pi(y)} - 1\Big| \le \varepsilon \right\}.$$

Remark 2.3. Note that we use a slightly different version of mixing time that the usual total-variation mixing. It turns out to be more convenient for our purposes.

Proof. We follow the *canonical paths* method by Diaconis and Saloff-Coste, see [19, Chapter 13].

Let D be the diagonal matrix with $D(x,x) = \sum_y Q(x,y)$. So

$$\mathbb{Q}_x[X_t=y] = e^{-t(D-Q)}(x,y).$$

Let Π be the diagonal matrix with π on the diagonal.

The main point is that because π is in detailed balance with Q, we get that $D-Q$ is self adjoint with respect to the inner product $\langle \Pi\cdot,\cdot\rangle$, which implies that there exists an orthonormal basis $\phi_1,\dots,\phi_{|S|}$ of eigenvectors for $D-Q$ with corresponding eigenvalues $\lambda_1 < \lambda_2 \le \dots \le \lambda_{|S|}$. Moreover, we have $\lambda_1 = 0$ and $\phi_1 = \vec{1}$.

Some short computations as in [19, Chapter 12] yield that for any x,

$$\sup_{x,y}\Big|\frac{\mathbb{Q}_x[X_t=y]}{\pi(y)} - 1\Big| \le e^{-t\gamma}\cdot\frac{1}{\min_x \pi(x)}.$$

Also, the min-max characterization (or Dirichlet form) gives

$$\lambda_2 = \min_{\substack{\varphi\neq\vec{0} \\ \langle\varphi,\pi\rangle=0}} \frac{\langle \Pi(D-Q)\varphi,\varphi\rangle}{\langle\Pi\varphi,\varphi\rangle}.$$

So, with the same proof as in the *comparison method* of [19, Chapter 13], an application of Cauchy-Schwarz gives that for any vector $\varphi\neq\vec{0}$ with $\langle\varphi,\pi\rangle = 0$,

$$\langle\Pi\varphi,\varphi\rangle \le \Theta\cdot\langle\Pi(D-Q)\varphi,\varphi\rangle.$$

Minimizing, we conclude that $\gamma \ge \Theta^{-1}$ which proves the lemma. ☐

Lemma 2.4. *Let* Q *be a matrix of rates for a continuous time Markov chain* (X_t) *on finite state space* S*, with measure* $\mathbb{Q}$*. Let* $T_x = \inf\{t\ge 0 \ : \ X_t = x\}$*. Then, for any* $t > \tau := 2\,t_{\mathrm{mix}}(1/2)$*, uniformly in* $x,y\in S$*,*

$$\mathbb{Q}_x[T_y > t] \le \exp\left(-\frac{t\pi(y)}{4\,\mathbb{E}_y[L(\tau,y)]}\right).$$

Proof. Fix $x, y \in S$. Note that for any $t > t_{\mathrm{mix}}(1/2)$ we have that

$$\mathbb{Q}_x[X_t = y] \geq \tfrac{\pi(y)}{2}.$$

We have

$$\mathbb{Q}_x[L(\tau, y) > 0] = \frac{\mathbb{E}_x[L(\tau, y)]}{\mathbb{E}_x[L(\tau, y) | L(\tau, y) > 0]} \geq \frac{\int_{t_{\mathrm{mix}}(1/2)}^{\tau} \mathbb{Q}_x[X_t = y] dt}{\mathbb{E}_y[L(\tau, y)]} \geq \frac{t_{\mathrm{mix}}(1/2)\pi(y)}{2\,\mathbb{E}_y[L(\tau, y)]}.$$

Note that $T_y > t$ if and only if $L(t, y) = 0$. Thus, because the above estimate is uniform in $x \in S$,

$$\mathbb{Q}_x[T_y > t] \leq \left(1 - \frac{t_{\mathrm{mix}}(1/2)\pi(y)}{2\,\mathbb{E}_y[L(\tau, y)]}\right)^{t/\tau} \leq \exp\left(-\frac{t\pi(y)}{4\,\mathbb{E}_y[L(\tau, y)]}\right),$$

as claimed. □

2.3 Flows and hitting estimates

Recall the assumptions (B1)–(B5). Given the measure π in these assumptions, we define the continuous time Markov chain $(X_t)_t$ on S, with law $\mathbb{Q}$, to be the chain whose transition rates are $Q(x, y) = \sqrt{\pi(y)/\pi(x)}\mathbf{1}_{\{x \sim y\}}$.

A few observations regarding this setup:

Lemma 2.5. *For the chain with law $\mathbb{Q}$ defined above, we have that for some constant $C > 0$ independent of L,*

$$t_{\mathrm{mix}}(1/2) \leq CL^2 \log L.$$

Proof. π is in detailed balance with Q.

For the collection of paths given in (B3), we have that for some constant $C > 0$ independent of L,

$$\begin{aligned}
\Theta = \Theta(Q, \pi) &= \max_{x \sim y} \frac{1}{\pi(x) Q(x, y)} \sum_{\substack{z, w \\ \gamma_{zw} \ni (x, y)}} \pi(z)\pi(w)|\gamma_{zw}| \\
&\leq \max_{x \sim y} \frac{(c_6)^2(\pi(x) \wedge \pi(y))^2}{\sqrt{\pi(x)\pi(y)}} \cdot L \cdot \#\{z, w \in S \ : \ (x, y) \in \gamma_{zw}\} \\
&\leq c_7 L^{d+2} \cdot \max_{x \sim y} \frac{(c_6)^2(\pi(x) \wedge \pi(y))^2}{\pi(x) \wedge \pi(y)} \\
&\leq c_7 (c_6)^2 L^{d+2} \max_x \pi(x) \leq CL^2,
\end{aligned}$$

where the last inequality follows from (B1). Using Lemma 2.2 and (B4) we conclude the proof. □

In order to bound the large deviations of this chain, we will find it useful to bound the expected local time accumulated up to time $\tau := 2\, t_{\mathrm{mix}}(1/2)$. This is the content of the next lemma. This lemma is also the main use of the *cone condition* (B2) above.

Lemma 2.6. *Let $\tau := 2\, t_{\mathrm{mix}}(1/2)$. For the chain with law $\mathbb{Q}$ defined above, we have that for some constant $C > 0$ independent of L, for any $y \in S$,*

$$\mathbb{E}_y[L(\tau, y)] \leq C(\log L)^3.$$

Proof. Fix $y \in S$. Let $\Sigma_m = \{z \in C_y \ : \ \mathrm{dist}(z, y) = m\}$ (the distance is the graph distance in $\mathbb{Z}^d$). Let $\Sigma = \Sigma_m$ for the largest m that is at most $\mathrm{diam}(C_z)$. Note that this m satisfies $m \geq c_4 L$ by (B2).

Step 1. First, we will bound from below the probability that the Markov chain started from y hits Σ before returning to y. Recall that our Markov chain with law $\mathbb{Q}$ is reversible. Therefore it is equivalent to a random walk on a network on S where the weight, or conductance, of the edge $e = (z, w)$ being given by $\pi(z)q(z, w) = \sqrt{\pi(z)\pi(w)}$. It is equivalent in the sense that both processes visit the same points in the same order, though possibly at different times. Hence it suffices to bound from above the effective resistance $\mathcal{R}_{\mathrm{eff}}(y \to \Sigma)$ in this network. Using (2.1) in (B2) above, $\pi(z) \geq c_5\pi(y)$ for all $z \in C_y$.

If we change the weights on all edges in C_y to be $c_5\pi(y)$ and the weights on edges not in C_y to be 0, then we have only reduced the conductance of all edges. Rayleigh's monotonicity principle (see [20, Chapter 2.4]) tells us that the effective resistance only increases. So it suffices to bound from above the effective resistance between y and Σ in this modified network.

The approach we use is that of [20, Chapter 2]. Let U be a random variable uniform on the base of the cone C_y at distance L from y. Let R be the ray from y to U. Let Γ be a (random) monotone path that stays at distance at most $\sqrt{2}$ from R, until reaching Σ. It is easily calculated that for an edge e at distance k from y in C_y, the probability that $e \in \Gamma$ is at most $O(k^{-(d-1)})$. Also, the number of edges in C_y at distance k from y is at most $O(k^{d-1})$. Finally, because Γ is chosen to be monotone, it traverses any edge at most once. So the function $\theta(e) = \mathbb{P}[e \in \Gamma] - \mathbb{P}[\hat{e} \in \Gamma]$ defines a unit flow in the cone C_y, from y to Σ (again see [20, Chapters 2.4 & 2.5]), where $\hat{e}$ is the directed edge e in the reverse direction. Thompson's principle ([20, Chapter 2.4]) then implies that the energy of this flow bounds the effective resistance:

$$\mathcal{R}_{\mathrm{eff}}(y \to \Sigma) \leq \sum_e \mathrm{res}(e)\theta(e)^2 \leq \sum_{k \leq m} \frac{1}{c_5\pi(y)} \cdot O(k^{-2(d-1)}) \cdot O(k^{d-1}) \leq \frac{1}{c_5\pi(y)} \cdot O(\log L).$$

Thus, if we define

$$T_A = \inf\{t \geq 0 \ : \ X_t \in A\},$$
$$T_y^+ = \inf\left\{t \geq T_{S\setminus\{y\}} \ : \ X_t = y\right\},$$

to be the hitting time of sets and return time to y, then for $w_y = \sum_{x \sim y} \sqrt{\pi(x)\pi(y)}$,

$$\mathbb{Q}_y[T_\Sigma < T_y^+] = \frac{1}{w_y\,\mathcal{R}_{\mathrm{eff}}(y \to \Sigma)} \geq (c \log L)^{-1}, \tag{2.8}$$

using (B5).

Step 2. We now provide a complementary bound for the tail of T_Σ. Let $(Y_n, n \geq 0)$ denote the jump chain of $(X_t, t \geq 0)$, and let $\mu(x) = \pi(x)q_x$ denote the invariant measure for Y; note that μ is then reversible for Y. Thus the Varopoulos-Carne bound ([20, Theorem 13.4]) states that

$$\mathbb{Q}_x[Y_n = z] \leq 2 \cdot \sqrt{\mu(z)/\mu(x)} \exp\left(-\frac{\mathrm{dist}(x, z)^2}{2n}\right).$$

Since $\mathrm{dist}(y,\Sigma) = m \geq cL$,

$$\mathbb{Q}_y[T_\Sigma(Y) < n] \leq \sum_{k=m}^{n} \mathbb{Q}_y[Y_k \in \Sigma]$$
$$\leq 2 \cdot \sum_{z\in\Sigma} \sqrt{\mu(z)/\mu(x)} \cdot \sum_{k=m}^{n} \exp(-cL^2/2k)$$

So with $n = \lfloor cL^2(\log L)^{-1} \rfloor$ for small enough constant $c > 0$,

$$\mathbb{Q}_y[T_\Sigma(Y) < n] \leq C \sum_{z\in\Sigma} \sqrt{\mu(z)/\mu(x)} \cdot \exp(-cL^2/n) = C \sum_{z\in\Sigma} \sqrt{\mu(z)/\mu(x)} \exp(-c^{-1}\log L).$$

On the other hand, using our assumptions (B1)–(B5) it is clear that $\sum_{z\in\Sigma} \sqrt{\mu(z)/\mu(x)} \leq CL^\alpha$ for some $\alpha > 0$. Thus choosing c small enough we have that

$$\mathbb{Q}_y[T_\Sigma(Y) \geq n] \to 1.$$

Since $c \leq q_x \leq c'$ for some $c, c' > 0$ and for all $x \in S$, we deduce that the total number of jumps of the jump chain by time t is at most a Poisson random variable with mean $c't$. Hence for some $c > 0$, for L large enough, by easy deviations for Poisson random variables,

$$\mathbb{Q}_y[T_\Sigma \geq cL^2(\log L)^{-1}] \geq 1/2. \tag{2.9}$$

Step 3. Let $H(s,t) = H_y(s,t)$ be the number of distinct visits to y in the time interval $[s,t)$; that is, the number of visits that its jump chain, restricted to jumps occurring in the interval $[s,t)$, pays to y during that time. We conclude from (2.8) that

$$\mathbb{E}_y[H(0,T_\Sigma)] = \frac{1}{\mathbb{Q}_y[T_\Sigma < T_y^+]} \leq c\log L. \tag{2.10}$$

We now use (2.10) to estimate $\mathbb{E}_y[L(\tau,y)]$. Define $S_0 = 0, T_0 = T_\Sigma$ and inductively,

$$S_j = \inf\{t \geq T_{j-1} \ : \ X_t = y\} \qquad \text{and} \qquad T_j = \inf\{t \geq S_j \ : \ X_t \in \Sigma\}.$$

These are the successive times for which the chain hits y then Σ and then y again, etc. The strong Markov property applied to (2.10) gives that for all j, $\mathbb{E}_y[H(S_j,S_{j+1})] \leq c\log L$.

Let $J = \max\{j \ : \ S_j \leq 2\, t_{\mathrm{mix}}(1/2)\}$. The events $(\{S_{j+1} - S_j \geq cL^2(\log L)^{-1}\})_j$ are independent, and each has probability at least $1/2$ by (2.9). So $\mathbb{E}_y[J] \leq c \cdot \frac{\tau\log L}{L^2}$. Thus,

$$\mathbb{E}_y[H(0,\tau)] \leq \sum_{j=1}^{\infty} \mathbb{E}_y[H(S_{j-1},S_j)\mathbf{1}_{\{S_{j-1}<\tau\}}] = \sum_{j=1}^{\infty} \mathbb{E}_y[H(S_{j-1},S_j)] \cdot \mathbb{Q}_y[S_{j-1} < \tau]$$
$$\leq c\log L \cdot \mathbb{E}_y[J] \leq c \cdot \frac{\tau(\log L)^2}{L^2}.$$

Together with Lemma 2.5 we have that

$$\mathbb{E}_y[H(0,\tau)] \leq c(\log L)^3.$$

Now, given that $H(0,\tau) = k$, the local time at y is at most the sum of k independent exponential random variables, each of mean $(\sum_z Q(y,z))^{-1} = \left(\sum_{z\sim y} \sqrt{\pi(z)/\pi(y)}\right)^{-1}$. Since there exists $z \in C_y$ with $Q(y,z) = \sqrt{\pi(z)/\pi(y)} \geq \sqrt{c_5}$ (by (2.1) in (B2)), we have that

$$\mathbb{E}_y[L(\tau,y)] \leq \mathbb{E}_y[H(0,\tau)] \cdot (c_5)^{-1/2} \leq c(\log L)^3,$$

as desired. ☐

We first derive the following corollary:

Corollary 2.7. *Let (X_t) be a Markov chain with transition rates given by Q. Let $T_y = \inf\{t \geq 0 : X_t = y\}$. Then uniformly over all $x \in S$,*

$$\mathbb{Q}_x(T_y > t) \leq \exp\left(-\frac{t\pi(y)}{c_{2.7}(\log L)^3}\right). \tag{2.11}$$

where $c_{2.7}$ is a uniform constant.

Proof. By Lemma 2.4, we have

$$\mathbb{Q}_x(T_y > t) \leq \exp\left(-\frac{t\pi(y)}{4\,\mathbb{E}_y(L(\tau,y))}\right).$$

By Lemma 2.6, $\mathbb{E}_y(L(\tau,y)) \leq C(\log L)^3$, so the result follows. ☐

2.4 Concentration estimates for local time

The next lemma is an auxiliary lemma regarding exponentially decaying random variables.

Lemma 2.8. *Suppose that T is a non-negative random variable with*

$$\mathbb{P}[T > t] \leq e^{-\mu t}$$

for all $t > 0$ and some $\mu > 0$. Then, for any $\alpha < 2^{-1/2}\mu$,

$$\mathbb{E}[e^{\alpha T}] \leq 1 + \alpha\,\mathbb{E}[T] + \frac{2(e+1)\alpha^2}{\mu^2}.$$

Proof. We use the inequality $e^x \leq 1 + x + x^2$ valid for all $x \leq 1$ (including negative x). This immediately implies the result for $\alpha \leq 0$. If $0 < \alpha < \mu$, then we have

$$\mathbb{E}[e^{\alpha T}] \leq 1 + \alpha\,\mathbb{E}[T] + \alpha^2\,\mathbb{E}[T^2] + \mathbb{E}[e^{\alpha T}\mathbf{1}_{\{\alpha T > 1\}}].$$

Clearly,

$$\mathbb{E}[T^2] \leq \int_0^\infty 2t\,\mathbb{P}[T > t]dt \leq \frac{2}{\mu^2}.$$

The last term can be bounded as follows:

$$\mathbb{E}[e^{\alpha T}\mathbf{1}_{\{\alpha T > 1\}}] = e\,\mathbb{P}[T > 1/\alpha] + \int_e^\infty \mathbb{P}[e^{\alpha T}\mathbf{1}_{\{\alpha T>1\}} > t]dt = e^{1-\frac{\mu}{\alpha}} + \int_{1/\alpha}^\infty \alpha e^{\alpha t}\,\mathbb{P}[T > t]dt$$
$$\leq e^{1-\frac{\mu}{\alpha}} + \alpha\int_{1/\alpha}^\infty e^{(\alpha-\mu)t}dt = (1 + \frac{\alpha}{\mu-\alpha}) \cdot e^{1-\frac{\mu}{\alpha}}.$$

Moreover, for any $x \geq \sqrt{2}$ we have that

$$(x-1)e^x \geq x^2 - 1 + \tfrac{x^3}{2} - \tfrac{x^2}{2} \geq \tfrac{x^3}{2}.$$

Thus, using $x = \mu/\alpha$, as long as $\sqrt{2}\alpha < \mu$, we have

$$\mathbb{E}[e^{\alpha T}] \leq 1 + \alpha\,\mathbb{E}[T] + 2x^{-2} + \frac{ex}{e^x(x-1)} \leq 1 + \alpha\,\mathbb{E}[T] + 2x^{-2} + 2ex^{-2}.$$

□

We now use the above results to deduce the following key step in the proof, which is an exponential concentration of the local time at a given point.

Lemma 2.9. *Let $(X_t)_t$ be a continuous time Markov chain on finite state space S with transition rates $Q(x,y) = \sqrt{\pi(y)/\pi(x)}\mathbf{1}_{\{x\sim y\}}$ for π satisfying assumption (B1)-(B5) above.*

For any $\delta > 0$, uniformly in $x, y \in S$,

$$\mathbb{Q}_x[L(t,y) \geq (1+\delta)\pi(y)t] \leq C\exp\left(-c_{2.9}\frac{\delta^2\pi(y)t}{(\log L)^6}\right) + C\exp\left(-c_{2.9}\delta\pi(y)t\right), \tag{2.12}$$

and for $0 < \delta < 1/2$,

$$\mathbb{Q}_x[L(t,y) \leq (1-\delta)\pi(y)t] \leq C\exp\left(-c_{2.9}\frac{\delta^2\pi(y)t}{(\log L)^6}\right). \tag{2.13}$$

Proof. Fix $y \in S$. In this proof it is convenient to define a time Δ by putting

$$\Delta = \frac{c_{2.7}(\log L)^3}{\pi(y)}. \tag{2.14}$$

($c_{2.7}$ is the constant from Corollary 2.7.) Intuitively, y is hit with positive probability (which can be made as high as desired) at least every Δ units of time by Corollary 2.7. Note that since $\pi(y) \leq c/L^d$, $\Delta \geq \tau = 2\,t_{\mathrm{mix}}(1/2)$ by Lemma 2.5.

We start with the proof of the lower bound on $L(t,y)$, (2.13), which is very similar except slightly harder than (2.12). Let $N(t)$ denote the number of distinct visit to y by time t. Let $q_y = \sum_{x\sim y} Q(y,x)$ be the total jump rate from y under $\mathbb{Q}$.

Let $n = \lceil \pi(y)q_y t(1-\delta)\rceil$. Note that by assumption (B5), $c \leq q_y \leq c'$ for some absolute constants c, c'. Moreover, if $L(t,y) \leq \pi(y)t(1-2\delta)$, then either $N(t) \leq n$ or $\sum_{i=1}^n E_i \leq \pi(y)t(1-2\delta) \leq \pi(y)t(1-\delta)^2 \leq (1-\delta)n/q_y$, where E_i are independent exponential random variables with rate q_y. The exponential moment of E_j is $\mathbb{E}[e^{-\alpha E_j}] = \frac{q_y}{q_y+\alpha}$ for all $\alpha > 0$. So an application of Chebyshev's inequality to the non-negative random variable $e^{-\alpha\sum_j E_j}$ gives

$$\mathbb{P}[\sum_{i=1}^n E_i \leq (1-\delta)n/q_y] \leq \left(\frac{q_y}{q_y+\alpha}\cdot e^{(1-\delta)/q_y}\right)^n.$$

Optimizing over α we obtain

$$\mathbb{P}[\sum_{i=1}^n E_i \leq (1-\delta)n/q_y] \leq \Big((1-\delta)e^{-\delta}\Big)^n = \exp\left(-c\delta\pi(y)t\right). \tag{2.15}$$

To bound the other term, it will be useful to define

$$T = \inf\{t \geq 0 \ : \ X_t = y\} \qquad T^+ = \inf\left\{t \geq T_{S\setminus\{y\}} \ : \ X_t = y\right\},$$

which are the hitting and return time to y, and also the successive return times to y: $T_0 = T$, and for $k > 0$,

$$S_k = \inf\{t \geq T_{k-1} \ : \ X_t \neq y\} \qquad T_k = \inf\{t \geq S_k \ : \ X_t = y\}.$$

Clearly $N(t) \leq n$ if and only if $T_n \geq t$. T_n is the sum of the independent increments $T_j - T_{j-1}$. By Corollary 2.7 each of these increments has an exponential tail $\mathbb{Q}_x[T_j > T_{j-1} + t] \leq e^{-\Delta^{-1}t}$. Thus, Lemma 2.8 (with $\mu = \Delta^{-1}$) gives that for any $0 < \alpha < \frac{1}{\sqrt{2}\Delta}$,

$$\mathbb{E}_x[e^{\alpha(T_j - T_{j-1})}] \leq 1 + \mathbb{E}_x[T_j - T_{j-1}] + 2(e+1)\alpha^2\Delta^2 \leq \exp\left(\alpha\,\mathbb{E}_x[T_j - T_{j-1}] + 2(e+1)\alpha^2\Delta^2\right).$$

Note that $\mathbb{E}_x[T_n] = n\,\mathbb{E}_y[T^+] + (\mathbb{E}_x[T^+] - \mathbb{E}_y[T^+])$. By Corollary 2.7, the second term is at most $\mathbb{E}_x[T^+] - \mathbb{E}_y[T^+] \leq \Delta$. Also, it is well known that $\mathbb{E}_y[T^+] = (q_y\pi(y))^{-1}$. So Chebyshev's inequality gives

$$\begin{aligned}\mathbb{Q}_x[N(t) \leq n] &= \mathbb{Q}_x(T_n \geq t) \leq \mathbb{E}_x(e^{\alpha T_n})e^{-\alpha t} \\ &\leq \exp\left(\alpha t(1-\delta) + \alpha\Delta + 2(e+1)\alpha^2\Delta^2(n+1) - \alpha t\right).\end{aligned}$$

Choosing $\alpha = \delta/(\pi(y)q_y\Delta^2)$, which is essentially optimal, we note that $0 < \alpha < \frac{1}{\sqrt{2}\Delta}$ for L sufficiently large, hence

$$\mathbb{Q}_x[N(t) \leq n] \leq C\exp\left(-c\frac{\delta^2\pi(y)t}{(\log L)^6}\right) \tag{2.16}$$

after simplification. Combining (2.15) and (2.16) gives (2.13).

The proof of (2.12) is similar. Here we choose $n = \lfloor \pi(y)q_yt(1+\delta)\rfloor$. If $L(t,y) \geq (1+\delta)^2\pi(y)t$ then either $N(t) \geq n$ or $\sum_{i=1}^n E_i \geq \pi(y)t(1+\delta)^2 \geq (n/q_y)(1+\delta)$. Simlarly to the argument above,

$$\mathbb{P}[\sum_{i=1}^n E_i \geq \pi(y)t(1+\delta)^2] \leq e^{-c\delta\pi(y)t}.$$

Also, by Lemma 2.8 and Corollary 2.7, as before,

$$\begin{aligned}\mathbb{Q}_x[N(t) \geq n] &\leq \mathbb{Q}_x[T_n \leq t] \leq \mathbb{E}[e^{-\alpha T_n}] \cdot e^{\alpha t} \\ &\leq C\exp\left(-\alpha t(1+\delta) + 2(e+1)\alpha^2\Delta^2 n + \alpha t\right) \\ &\leq C\exp\left(-c\frac{\delta^2\pi(y)t}{(\log L)^6}\right).\end{aligned}$$

This completes the proof, continuing as in the lower bound. It is important here that we do not restrict ourselves to $\delta < 1$, so for large δ, the bound $e^{-c\delta\pi(y)t}$ kicks in. □

2.5 Change of measure

The following analogue of the Gauss-Green formula will be useful.

Lemma 2.10. *Let ϕ, ψ be two given functions with support contained in a finite set S. For any $D \subset S$,*

$$\sum_{x\in D} \phi(x)\mathcal{L}\psi(x) = -\tfrac{1}{2}\mathcal{E}_D(\phi,\psi) + \sum_{x\in D, y\notin D : x\sim y} \phi(x)\nabla\psi(x,y). \tag{2.17}$$

(This is the analogue of the classical Gauss-Green formula $\int f\Delta g = -\int \nabla f \nabla g$ valid for smooth functions f, g with compact support, but note the additional boundary terms.)

Proof. We have

$$\begin{aligned}\mathcal{E}_D(\phi,\psi) &= \sum_{x\sim y\in D} \nabla\phi(x,y)\nabla\psi(x,y) = \sum_{x\sim y\in D}\phi(y)\nabla\psi(x,y) - \sum_{x\sim y\in D}\phi(x)\nabla\psi(x,y) \\ &= -2\sum_{x\in D}\phi(x)\mathcal{L}\psi(x) + 2\sum_{x\in D}\sum_{D\not\ni y\sim x}\phi(x)\nabla\psi(x,y).\end{aligned}$$

□

The crucial change of measure calculation is provided by the following lemma.

Lemma 2.11. *Let π be some probability vector and let S be a connected subset of $\mathbb{Z}^d$ containing the support of π, $\{\pi > 0\} \subset S$. Let $\mathbb{Q}$ be the law of the continuous time Markov chain with transition rates $Q(x,y) = \sqrt{\pi(y)/\pi(x)}\mathbf{1}_{\{x\sim y\}}$.*

Let Z_t denote the Radon-Nikodym derivative of $\mathbb{P}$ with respect to $\mathbb{Q}$, restricted to $\mathcal{F}_t$:

$$Z_t := \frac{d\mathbb{P}}{d\mathbb{Q}}\Big|_{\mathcal{F}_t}. \tag{2.18}$$

Define

$$S_- = \{x \in S \ : \ \mathcal{L}f(x) \le 0\} \qquad \text{and} \qquad S_+ = \{x\in S \ : \ \mathcal{L}f(x) > 0\}$$

is the domains where f is respectively super-harmonic or sub-harmonic.

Then, for any

$$\omega \in \mathcal{D}_{\alpha,\beta} = \{\alpha\pi(x) \le \frac{L(t,x)}{t} \le \beta\pi(x), x\in\mathbb{Z}^d\},$$

we have the pointwise bounds:

$$Z_t(\omega) \ge \sqrt{\frac{\min\pi}{\max\pi}}\exp\left(-t\cdot\tfrac{\beta}{2}\cdot\mathcal{E}(f,f)\right)\cdot\exp\left(-t\cdot(\beta-\alpha)\cdot\Phi(f)\right), \tag{2.19}$$

and likewise,

$$Z_t(\omega) \le \sqrt{\frac{\max\pi}{\min\pi}}\exp\left(-t\cdot\tfrac{\alpha}{2}\mathcal{E}(f,f)\right)\cdot\exp\left(t\cdot(\beta-\alpha)\cdot\Phi(f)\right). \tag{2.20}$$

where $f = \sqrt{\pi}$ and

$$\Phi(f) = \sum_{\substack{x\in S_+, y\in S_-\\ x\sim y}} f(x)\nabla f(x,y).$$

Specifically, if f is sub-harmonic, then $\Phi(f) = 0$ and

$$\sqrt{\frac{\min\pi}{\max\pi}}\cdot\exp\left(-t\cdot\tfrac{\beta}{2}\mathcal{E}(f,f)\right) \le D_t(\omega) \le \sqrt{\frac{\max\pi}{\min\pi}}\cdot\exp\left(-t\cdot\tfrac{\alpha}{2}\mathcal{E}(f,f)\right).$$

Proof. Let us start by expanding the energy $\mathcal{E}(f,f)$. Because f vanishes off S, the Gauss-Green formula (Lemma 2.10) gives

$$\mathcal{E}(f,f) = -2\sum_{x\in S} f(x)\mathcal{L}f(x) = -2\sum_{x\in S_+} f(x)\mathcal{L}f(x) - 2\sum_{x\in S_-} f(x)\mathcal{L}f(x).$$

Let

$$B := -\sum_{x\in S_-} f(x)\mathcal{L}f(x) \qquad \text{and} \qquad A := \sum_{x\in S_+} f(x)\mathcal{L}f(x).$$

So $B, A > 0$ and $B - A = \frac{1}{2}\mathcal{E}(f,f)$.

Also, again by Lemma 2.10,

$$A = -\frac{1}{2}\mathcal{E}_{S_+}(f,f) + \sum_{\substack{x\in S_+, y\in S_- \\ x\sim y}} f(x)\nabla f(x,y) - \sum_{\substack{x\in S_+, y\notin S \\ x\sim y}} f(x)^2$$

and similarly,

$$B = \frac{1}{2}\mathcal{E}_{S_-}(f,f) - \sum_{\substack{x\in S_-, y\in S_+ \\ x\sim y}} f(x)\nabla f(x,y) + \sum_{\substack{x\in S_-, y\notin S \\ x\sim y}} f(x)^2.$$

Note that

$$\frac{1}{2}\mathcal{E}(f,f) = \frac{1}{2}\mathcal{E}_{S_-}(f,f) + \frac{1}{2}\mathcal{E}_{S_+}(f,f) + \sum_{\substack{x\in S_+, y\in S_- \\ x\sim y}} (f(y)-f(x))^2 + \sum_{\substack{x\in S, y\notin S \\ x\sim y}} f(x)^2.$$

Thus, since $\mathcal{E}_D(f,f) \geq 0$, we can bound

$$\begin{aligned}
B + A &= \frac{1}{2}\left(\mathcal{E}_{S_-}(f,f) - \mathcal{E}_{S_+}(f,f)\right) + \sum_{\substack{x\in S_+, y\in S_- \\ x\sim y}} (f(x)+f(y))\nabla f(x,y) + \sum_{\substack{x\in S_-, y\notin S \\ x\sim y}} f(x)^2 - \sum_{\substack{x\in S_+, y\notin S \\ x\sim y}} f(x)^2 \\
&\leq \frac{1}{2}\left(\mathcal{E}_{S_-}(f,f) + \mathcal{E}_{S+}(f,f)\right) + \sum_{\substack{x\in S, y\notin S \\ x\sim y}} |\nabla f(x,y)|^2 + \sum_{\substack{x\in S_+, y\in S_- \\ x\sim y}} (f(y)^2 - f(x)^2) \\
&= \frac{1}{2}\sum_{x\sim y} |\nabla f(x,y)|^2 - \sum_{\substack{x\in S_+, y\in S_- \\ x\sim y}} (f(y)-f(x))^2 + \sum_{\substack{x\in S_+, y\in S_- \\ x\sim y}} (f(y)^2 - f(x)^2) \\
&= \frac{1}{2}\mathcal{E}(f,f) + 2\sum_{\substack{x\in S_+, y\in S_- \\ x\sim y}} f(x)\nabla f(x,y) = \frac{1}{2}\mathcal{E}(f,f) + 2\Phi(f).
\end{aligned}$$

Since $B > A$, this implies that for any $0 < \alpha < \beta$,

$$-\beta B + \alpha A = -\frac{\beta+\alpha}{2}(B-A) - \frac{\beta-\alpha}{2}(A+B) \geq -\frac{\beta}{2}\mathcal{E}(f,f) - (\beta-\alpha)\Phi(f). \tag{2.21}$$

Similarly,

$$-\alpha B + \beta A = -\frac{\beta+\alpha}{2}(B-A) + \frac{\beta-\alpha}{2}(A+B) \leq -\frac{\alpha}{2}\mathcal{E}(f,f) + (\beta-\alpha)\Phi(f). \tag{2.22}$$

Let $\omega \in \mathcal{D}_{\alpha,\beta}$. Suppose the successive states visited by ω up to time t are $x_0, \ldots, x_n$, with the path staying a time $\tau_0, \ldots, \tau_n$ at respectively at these locations. (Hence $\tau_1 + \ldots + \tau_n = t$.) Assume $\omega \in \mathcal{D}_{\alpha,\beta}$. If $x \in \mathbb{Z}^d$, then the total rate at which the particle would jump out of x under $\mathbb{Q}$ is given by $q(x) = f(x)^{-1} \sum_{y \sim x} f(y)$.

Then a quick calculation shows that, letting $d(x) = 2d$ be the total rate of leaving x under $\mathbb{P}$,

$$\begin{aligned}\frac{d\mathbb{P}}{d\mathbb{Q}}(\omega) &= \frac{e^{-d(x_0)\tau_0} \ldots e^{-d(x_n)\tau_n}}{Q(x_0, x_1)e^{-q(x_0)\tau_0} \ldots Q(x_{n-1}, x_n)e^{-q(x_n)\tau_n}} \\ &= \frac{f(x_n)}{f(x_0)} \prod_{i=0}^{n} \exp\{(q-d)(x_i)\tau_i\} \\ &= \frac{f(x_n)}{f(x_0)} \prod_{x \in \mathbb{Z}^d} \exp\{(q-d)(x)L(t,x)\} \\ &= \frac{f(x_n)}{f(x_0)} \exp\left\{ \sum_{x \in \mathbb{Z}^d} \frac{\mathcal{L}f(x)}{f(x)} L(t,x) \right\}.\end{aligned}$$

We split this sum into a sum on S_- and S_+. Using the fact that $\omega \in \mathcal{D}_{\alpha,\beta}$, (2.21) provides the lower bound

$$\frac{d\mathbb{P}}{d\mathbb{Q}}(\omega) \geq \exp\left(t \cdot (-\beta B + \alpha A)\right) \geq \exp\left(-t \cdot \tfrac{\beta}{2}\mathcal{E}(f,f)\right) \cdot \exp\left(-t \cdot (\beta - \alpha) \cdot \Phi(f)\right),$$

which proves (2.19). Proving (2.19) from (2.22) is similar. □

This section is concluded by

Proof of Theorem 2.1. The upper bound is a direct application of the upper bound in Lemma 2.11. For the lower bound, using the lower bound in Lemma 2.11, we need to show that

$$\mathbb{Q}[\mathcal{D}_{\alpha,\beta}] \geq \sigma = 1 - C \sum_{y \in S} \exp\left(-c(\alpha - 1)^2 \pi(y) t \cdot (\log L)^{-6}\right).$$

This follows from (2.12) and (2.13) with suitable choice of δ. □

3 Proof of Theorem 1.1

3.1 Partition function estimate

The goal of this section is to obtain the following lower bound on the partition function.

Proposition 3.1.

$$Z_{t,\beta} \geq \exp(-c(\log t)^5 \cdot t^{1/3})$$

for some constant c depending only on β.

Remark 3.2. *Our proof relies on Theorem 2.1. Ofer Zeitouni has pointed out to us that there is another and perhaps altogether simpler route for obtaining this lower bound using the methods developed in [6]. We have chosen to keep the proof as it was because we believe that Theorem 2.1 may prove useful for obtaining finer results than the ones we discuss here.*

Let L be the nearest odd integer to $t^{1/3}$. To prove Proposition 3.1 we introduce a square S of side length $2L+1$ and containing the origin 0 as an arbitrary fixed point (*e.g.* , the center); that is, $S = [-L, L]^2 \cap \mathbb{Z}^2$.

The argument relies on an appropriate change of measure and uses the techniques developed in the preceding section. We proceed to define the local time profile π. For $z \in S$ define

$$\pi(z) = \frac{\operatorname{dist}(z, \partial S)}{L^3}, \tag{3.1}$$

and $\pi(z) = 0$ for $z \notin S$. Let $f = \sqrt{\pi}$. Note that because $2\sqrt{a} \geq \sqrt{a+1} + \sqrt{a-1}$ for $a > 0$, we have that

$$\mathcal{L}f(z) \leq 0 \ , \qquad \forall\ z \in S. \tag{3.2}$$

That is, f is sub-harmonic. It is also not difficult to verify that π satisfies (B1)-(B5). Let $\mathbb{Q}$ be the law of the continuous time Markov chain on S with transition rates given by $Q(x,y) = \sqrt{\pi(y)/\pi(x)}\mathbf{1}_{\{x\sim y\}}$.

Let $K = K(L) = C(\log L)^4$ for some large enough constant C (independent of L, t) to be chosen below. We define the event

$$\mathcal{A} := \{\forall z \in S, L(t,z) < K\pi(z)t\}. \tag{3.3}$$

We also define the event $\mathcal{B}$:

$$\mathcal{B} := \{|\partial R_t| \leq \ CL(\log L)^3\}.$$

Finally, define the good event

$$\mathcal{G} = \mathcal{A} \cap \mathcal{B}. \tag{3.4}$$

Proposition 3.1 will follow immediately from the following estimate:

$$\mathbb{P}[\mathcal{B}] \geq \mathbb{P}[\mathcal{G}] \geq \exp(-c(\log t)^5 t^{1/3}), \tag{3.5}$$

since for any $\omega \in \mathcal{B}$, $|\partial R_t| \leq CL(\log L)^3$ and hence the energy satisfies $H(\omega) \geq \exp(-cL(\log L)^3) \geq \exp(-ct^{1/3}(\log t)^3)$.

We will prove (3.5) by insisting that the random walk stay in S with a certain well-defined local time profile π within S.

Remark 3.3. *Note that it is quite simple to estimate the probability to stay in the square S for time t, see e.g. Lemma 3.7 later on, which gives a slightly better lower bound than Proposition 3.1. However if we simply condition on staying in S, the actual boundary of the range might in fact be much larger than L, the boundary of the square S. This is because conditionally on staying in S, points near the boundary will in fact not be visited, as a quick calculation shows. In fact, the boundary of the range will then be typically at distance $\sqrt{L}$ from the boundary of S, and probably is a very irregular (fractal) curve.*

We turn back to the event $\mathcal{G}$ defined in (3.4). The following proposition says that the good event $\mathcal{G}$ has a reasonable probability under the measure $\mathbb{Q}$.

Proposition 3.4. *For L sufficiently large, $\mathbb{Q}[\mathcal{G}] \geq 1/4$.*

Proof. Using Lemma 2.9, and specifically (2.12), for any $z, w \in S$,

$$\mathbb{Q}_z[L(t,w) \geq K\pi(w)t] \leq \exp\left(-c(\log L)^2\pi(w)t\right),$$

where the constant $c > 0$ can be made large by changing C in the definition of $K(L)$. As long as $c\pi(w)t \geq 1$ for all $w \in S$, a union bound on $w \in S$ gives that $\mathbb{Q}_z[\mathcal{A}] = 1 - O(L^2 \cdot e^{-(\log L)^2})$ as $L \to \infty$.

Now, we estimate $\mathbb{E}[|\partial R_t|]$ under $\mathbb{Q}$. Since the maximal degree is 4 we obtain that $|\partial R_t| \leq 4|S \setminus R_t|$. Using Corollary 2.7

$$\mathbb{E}[|S \setminus R_t|] = \sum_{z \in S} \mathbb{Q}[z \notin R_t] = \sum_{z \in S} \mathbb{Q}[T_z > t] \leq \sum_{z \in S} \exp\left(-c\frac{\pi(z)t}{(\log L)^3}\right).$$

For $1 \leq r \leq L$, the vertices in S of distance r from ∂S are exactly $\{-(L-r), L-r\} \times [-(L-r), L-r] \bigcup [-(L-r), L-r] \times \{-(L-r), L-r\}$, which is the square boundary of side-length $2(L-r)+1$. This set is of size $8(L-r)$ (except for $r = L$ where it is of size 1). So, with $\Lambda = c\frac{t}{L^3(\log L)^3}$,

$$\begin{aligned}\mathbb{E}[|S \setminus R_t|] &\leq e^{-L\Lambda} + \sum_{r=1}^{L-1} 8(L-r)e^{-r\Lambda} = e^{-L\Lambda} + 8e^{-L\Lambda} \cdot \sum_{x=1}^{L-1} e^{x\Lambda} \\ &\leq e^{-L\Lambda} \cdot \left(1 + 8 \cdot \int_1^L xe^{x\Lambda}dx\right) = O(L^4(\log L)^3 \cdot t^{-1}).\end{aligned}$$

Thus, by our choice of $L = t^{1/3}$,

$$\mathbb{Q}_z[|\partial R_t| \geq CL(\log L)^3] \leq (c'/C)L^3t^{-1} = (c'/C).$$

This is at most $1/2$ if $C > 0$ is large enough.

Both estimates together give

$$\mathbb{Q}_z[\mathcal{G}] \geq \mathbb{Q}_z[\mathcal{B}] - \mathbb{Q}_z[\mathcal{A}^c] \geq \tfrac{1}{2} - o(1),$$

as $L \to \infty$. □

We now turn to:

Proof of Proposition 3.1. By Lemma 2.11 and Proposition 3.4, we have that

$$\mathbb{P}(\mathcal{G}) \geq \frac{c}{L}\exp(-Kt\mathcal{E}(f,f)).$$

Thus (3.5), and hence Proposition 3.1, will follow from the estimate $\mathcal{E}(f,f) \leq c\log L \cdot L^{-2}$. For $1 \leq r \leq L$ let $S_r = \{-(L-r), L-r\} \times [-(L-r), L-r] \bigcup [-(L-r), L-r] \times \{-(L-r), L-r\}$ be the set of vertices which are at distance r from ∂S. This set is of size $8(L-r)$ if $r \leq L-1$ and 1 if $r = L$. For $1 \leq r < L$, if $z \sim w$ for $z \in S_{r+1}, w \in S_r$ then a first order Taylor expansion gives

$$|f(z) - f(w)| \leq L^{-3/2} \cdot \frac{1}{2\sqrt{r}}.$$

If $z \in S_1$ and $w \in S_0$ then $|f(z) - f(w)| = L^{-3/2}$. Thus,

$$\begin{aligned}\mathcal{E}(f,f) &\leq \sum_{r=0}^{L-1} \sum_{z \in S_{r+1}, w \in S_r} |f(z) - f(w)|^2 \\ &\leq L^{-3} \cdot \sum_{r=1}^{L-2} \frac{1}{4r} \cdot 8(L-r-1) + L^{-3} 4 \cdot \frac{1}{4(L-1)} + 8(L-1)L^{-3} \\ &\leq O(L^{-2} \log L)\end{aligned}$$

□

3.2 Discrete planar isoperimetry

We now state and prove a modified isoperimetric inequality which deals with the outer boundary of a set. We first need some definitions. For a set $G \subset \mathbb{Z}^2$, let $\mathrm{Ext}(G)$ be the unique unbounded connected component of $\mathbb{Z}^2 \setminus G$. Let the outer vertex boundary $\partial^* G$ be defined by

$$\partial^* G = \{x \in G : \exists y \in \mathrm{Ext}(G), x \sim y\}.$$

The outer edge boundary, denoted by $\partial_e^* G$, consists of those edges $e = (x,y)$ with $x \in G$ and $y \in \mathrm{Ext}(G)$.

Lemma 3.5. *Let $G \in \mathcal{G}$. Let R be the smallest rectangle in $\mathbb{Z}^2$ containing G. Then, $|\partial^* R| \leq 3|\partial^* G|$. Furthermore, $|G| \leq |R| \leq |\partial^* G|^2$.*

Proof. For any connected set S, we have that $|\partial^* S| \leq |\partial_e^* S| \leq 3|\partial^* S|$ since adjacent to any vertex in $\partial^* S$ there are at most 3 edges in $\partial_e^* S$ (and at least one edge must connect that vertex to the set itself). Thus, it suffices to prove that $|\partial_e^* R| \leq |\partial_e^* G|$.

Let $(x, x+e) \in \partial_e^* G$, for some $e \in \{(\pm 1, 0), (0, \pm 1)\}$. Then $x \in G$ and hence $x \in R$. Thus, there exists a (necessarily unique) $k \geq 0$ such that $x + ke \in R$ and $x + ze \notin R$ for all $z > k$. Thus, $(x + ke, x + (k+1)e) \in \partial_e^* R$. Define a map $\phi : \partial_e^* G \to \partial_e^* R$ by $\phi((x, x+e)) = (x + ke, x + (k+1)e)$ for this k.

We claim that $\phi : \partial_e^* G \to \partial_e^* R$ is onto. This follows since if $(x, x+e) \in \partial_e^* R$, then considering the line $L = \{x - ke \ : \ k \geq 0\}$, it must be that $L \cap G \neq \emptyset$, since otherwise either G would not be connected or R would not be the smallest rectangle containing G. Thus, there must exist some $k \geq 0$ such that $x - ke \in G$ and $x - ze \notin G$ for any $z < k$. Thus, the edge $(x - ke, x - (k-1)e)$ is in $\partial_e^* G$, and it is immediate that $\phi(x - ke, x - (k-1)e) = (x, x+e)$.

This proves that there is a map from $\partial_e^* G$ onto $\partial_e^* R$. So $|\partial^* R| \leq |\partial_e^* R| \leq |\partial_e^* G| \leq 3|\partial^* G|$.

The second assertion follows from the fact that $(k+m)^2 \geq 4km$. So for a $k \times m$ rectangle R, $|\partial R|^2 = 4(k+m)^2 \geq 16km = 16|R|$. □

3.3 Proof of condensation

We will prove the following more precise statement of Theorem 1.1.

Theorem 3.6. *Assume $\beta > \beta_0 = \log \alpha$, where α is the connective constant. There exist constants $c_1, c_2 > 0$ depending on β only such that the following holds:*

$$\mu_t \Big[c_1 t^{1/3} (\log t)^{-5} \leq \mathrm{diam}(R_t) \leq c_2 t^{1/3} (\log t)^5 \Big] \geq 1 - \exp\Big(-c t^{1/3} (\log t)^5 \Big). \tag{3.6}$$

With these preliminaries we are now ready to prove Theorem 3.6.

Proof of Theorem 3.6. Let L be the closest integer to $t^{1/3}$. Recall that by Proposition 3.1

$$Z(t,\beta) \geq \exp(-ct^{1/3}(\log t)^5) \tag{3.7}$$

where $c > 0$ is a constant depending only on β.

Now for $k \geq 1$, let Γ_k be the set of closed simple loops $\gamma : \{0, \dots, k\} \to \mathbb{Z}^2$ of length k (and thus $\gamma(0) = \gamma(k)$. We will abuse notations and also write γ for the trace of γ, i.e., the set $\{\gamma(0), \dots, \gamma(k-1)\}$. For $\gamma \in \Gamma_k$, we will estimate the probability that $\partial^* R_t = \gamma$. Note that

$$\limsup_{k\to\infty} |\Gamma_k|^{1/k} \leq \alpha, \tag{3.8}$$

where α is the connective constant of $\mathbb{Z}^2$. Let $\beta_0 = \log \alpha$ and choose $\beta > \beta_0$. Choose also $\beta' = \frac{\beta+\beta_0}{2}$ so that $\beta_0 < \beta' < \beta$.

Let $M > 0$ be large enough so that $|\Gamma_j| \leq e^{\beta' j}$ for all $j \geq M$. Then,

$$\begin{aligned} \mu_t[|\partial^* R_t| > M] &\leq \sum_{j=M}^{t} \sum_{\gamma \in \Gamma_j} e^{-\beta j} Z(t,\beta)^{-1}\, \mathbb{P}[\partial^* R_t = \gamma] \\ &\leq \sum_{j=M}^{\infty} e^{\beta' j - \beta j} \exp(ct^{1/3}(\log t)^5) \\ &\leq \left(1 - e^{-\frac{1}{2}(\beta-\beta_0)M}\right)^{-1} \cdot \exp\left(ct^{1/3}(\log t)^5 - \tfrac{1}{2}(\beta-\beta_0)M\right). \end{aligned}$$

Thus, if we take

$$M \geq C\tfrac{c}{\beta-\beta_0} t^{1/3}(\log t)^5, \tag{3.9}$$

with $C \geq 4$, we get

$$\mu_t[|\partial^* R_t| > M] \leq \exp(-ct^{1/3}(\log t)^5). \tag{3.10}$$

Now, it is an obvious geometric fact that for any shape $G \subset \mathbb{Z}^2$, if R is the smallest rectangle that contains it then

$$\operatorname{diam}(G) \leq |\partial^* R| \leq 3|\partial^* G|, \tag{3.11}$$

where the second inequality comes from Lemma 3.5. We deduce that

$$\mu_t[\operatorname{diam}(R_t) > M] \leq \exp(-ct^{1/3}(\log t)^5).$$

Since $\operatorname{diam}(R_t) \leq M$ implies that $R_t \subset [-M, M]^2$, we have that

$$\mu_t[|R_t| > (2M+1)^2] \leq \exp(-ct^{1/3}(\log t)^5).$$

This bounds from above the external boundary size $|\partial^* R_t|$, diameter $\operatorname{diam}(R_t)$ and volume $|R_t|$.

For the lower bound on the diameter and boundary size we require the following estimate.

Lemma 3.7. *Let R be a rectangle of sidelengths J and K respectively. Then if $z \in R$,*

$$\mathbb{P}_z[R_t \subset R] \leq JK \exp\left(-t\tfrac{\pi^2}{4}(\tfrac{1}{J^2}+\tfrac{1}{K^2})\right) \leq JK \exp\left(-\frac{\pi^2 t}{2|R|}\right).$$

Proof. For the purpose of this proof we may assume (by standard deviation bounds on Poisson random variables) that $\mathbb{P}$ is the law of continuous time simple random walk on $\mathbb{Z}^2$. In this case the coordinates X_t, Y_t are independent rate $(1/2)$ simple random walks on $\mathbb{Z}$. We just focus on X_t and compute $\mathbb{P}_x(T > t)$ where $x \in \{1, \dots, J\}$ and $T = \inf\{t \geq 0 : X_t \notin [1, J]\}$. Let $\mathcal{L}$ denote the generator of (rate 1) simple random walk on $\mathbb{Z}$, and let $\phi(x) = e^{i\pi x/J}$. It is trivial to check that

$$\mathcal{L}\phi(x) = -\lambda\phi(x)$$

for all $x \in \mathbb{Z}$, where $\lambda = 1 - \cos(\pi/J)$. Thus if we let $\psi(t, x) = e^{\lambda t}\sin(\pi x/J)$ we have

$$\frac{\partial}{\partial t}\psi + \mathcal{L}\psi = 0$$

and hence $M_t := e^{\lambda t}\sin(\pi X_t/J)$ is a martingale. Consequently, applying the optional stopping time theorem at the time $t \wedge T$ (which is bounded), and the inequality $\sin(u) \geq (2/\pi)u$ valid for $0 \leq u \leq \pi/2$, yields

$$\begin{aligned}\sin(\pi x/J) &= \mathbb{E}_x(e^{\lambda t}\sin(\pi X_{t\wedge T}/J)) \\ &\geq e^{\lambda t}\frac{2}{J}\,\mathbb{P}_x(T > t).\end{aligned}$$

Therefore,

$$\mathbb{P}_x(T > t) \leq Je^{-\lambda t}.$$

Now, $\lambda = 1 - \cos(\pi/J) \geq \pi^2/4J^2$ for J large enough, and the result follows.

The last inequality is just $2JK \leq J^2 + K^2$ which always holds. □

We now combine Lemma 3.7 with (3.10) to obtain a lower bound on the diameter of R_t under μ_t:

Let $R_t^\square$ be smallest rectangle containing R_t. Let J_t, K_t be the side-lengths of $R_t^\square$. Lemma 3.5 tells us that

$$2(J_t + K_t) - 4 = |\partial R_t^\square| = |\partial^* R_t^\square| \leq 3|\partial^* R_t|.$$

Note also that

$$|R_t^\square| = J_t K_t \leq \left(\tfrac{J_t+K_t}{2}\right)^2 \leq \tfrac{1}{4}\left(\tfrac{3|\partial^* R_t|+4}{2}\right)^2.$$

If $J_t = j, K_t = k$ then $R_t \subset [x, x+k] \times [y, y+j]$ for some $|x| \leq k, |y| \leq j$. Also, by Lemma 3.5,

$$|\partial R_t| \geq |\partial^* R_t| \geq \frac{2(J_t + K_t) - 4}{3}.$$

So by Lemma 3.7, for an integer $N > 0$,

$$\begin{aligned}\mu_t[J_t, K_t \leq N] &\leq \sum_{k,j=1}^{N} \mu_t[K_t = k, J_t = j] \\ &\leq Z(t,\beta)^{-1} \cdot \sum_{k,j=1}^{N} \exp\left(-\beta \frac{2(k+j)-4}{3}\right) \sum_{x=-k}^{k} \sum_{y=-j}^{j} \mathbb{P}[R_t \subset [x, x+k] \times [y, y+j]] \\ &\leq Z(t,\beta)^{-1} \cdot \sum_{k,j=1}^{N} \exp\left(-\beta \frac{2(k+j)-4}{3}\right) \cdot (2k+1)(2j+1)jk \exp\left(-\frac{\pi^2 t}{2kj}\right) \\ &\leq Z(t,\beta)^{-1} C' \exp\left(-\frac{\pi^2 t}{2N^2}\right). = C' \exp\left(ct^{1/3}(\log t)^5 - \frac{\pi^2}{2} \cdot \frac{t}{N^2}\right).\end{aligned}$$

Thus, if

$$N \leq \frac{\pi^2}{4c} t^{1/3} (\log t)^{-5},$$

we get that

$$\mu_t[J_t, K_t \leq N] \leq \exp\left(-ct^{1/3}(\log t)^5\right).$$

Since these are the dimensions of the smallest rectangle containing R_t, with high μ_t-probability

$$|\partial R_t| \geq c'N \qquad \text{and} \qquad \operatorname{diam}(R_t) \geq c'N,$$

for some small enough $c' > 0$. $\square$

3.4 Proof of Theorems 1.3 and 1.4

We explain how to adapt the arguments of the proof of Theorem 1.1 to give the proof of Theorem 1.4. Let $T_y(\beta) = \inf\{t \geq 0 : L(t,y) \geq \beta\}$. The argument of Lemma 2.4 shows that

$$\mathbb{P}(T_y(\beta) > t) \leq \exp\left(-\frac{t\pi(y)}{4\,\mathbb{E}_y(L(\tau,y))} \cdot \frac{\tau}{\tau+\beta} e^{-\beta q_y}\right) \leq \exp\left(-c\frac{t\pi(y)}{\mathbb{E}_y(L(\tau,y))}\right).$$

Thus by Lemma 2.6,

$$\mathbb{P}(T_y(\beta) > t) \leq \exp\left(-\frac{t\pi(y)}{c(\log L)^3}\right),$$

which is the analogue of Corollary 2.7. Now, recall the events $\mathcal{A}$ in the proof of Proposition 3.1:

$$\mathcal{A} = \{\forall z \in S, L(t,z) \leq K\pi(z)t\};$$

We modify the definition of $\mathcal{B}$. Let $S' = \{z \in S : \operatorname{dist}(z, \partial S) \geq (\log L)^4\}$ and consider the event

$$\mathcal{B}' = \{\forall z \in S', L(t,z) \geq \beta\}.$$

Let $\mathcal{G}' = \mathcal{G}'_t = \mathcal{A} \cap \mathcal{B}'$. Then $\mathbb{Q}(\mathcal{G}') \geq 1/4$ for L sufficiently large. Indeed . Then

$$\mathbb{Q}(\mathcal{B}'^{\complement}) \leq \sum_{z \in S'} \mathbb{Q}(T_z(\beta) > t) \leq 1/2$$

for L large enough, exactly as in the proof of Proposition 3.4. We deduce that

$$\mathbb{P}(\mathcal{G}') \geq \exp(-c(\log t)^5 t^{1/3}).$$

Assume that $\mathcal{G}'_{t/2}$ holds. In the remaining $t/2$ units of time, we make sure that the each of the remaining $O(L(\log L)^4)$ vertices of $S \setminus S'$ are visited at least β units of time each, as follows: for each $1 \leq k \leq (\log L)^4$ let $S_k = \{z \in S : \mathrm{dist}(z, \partial S) = k\}$. We visit each vertex in S_k in clockwise order, starting from $(-L+k, 0)$. At each new vertex, the walk remains at least β and at most 2β units of time. When the walk has visited each vertex of S_k, it moves on to S_{k+1}. The total amount of time spend doing so is at most $2\beta L(\log L)^4$, which is much less that $t/2$. In the remaining amount of time, the walk is free to do what it wants, provided it stays in S. If all these conditions are fulfilled, it is clear that $R_t = S$ and that each vertex has a local time greater than β, so $\mathcal{E}_t$ holds. The probability of visiting every vertex in this prescribed order immediately after $t/2$ is at least $\exp(-cL(\log L)^4)$ for some $c = c(\beta) < \infty$. The probability of remaining in S after that is at least $\exp(-K(t/2)\mathcal{E}(f, f))$ and hence at least $\exp(-c(\log t)^5 t^{1/3})$. All in all, we deduce

$$\mathbb{P}(\mathcal{E}_t) \geq \exp(-c(\log t)^5 t^{1/3}). \tag{3.12}$$

This argument also shows that if $\tilde{Z}(t, \beta)$ is the partition function corresponding to the Hamiltonian $\tilde{H} = \sum_{x \in \partial R_t} L(t, x)$ in (1.4), then

$$\tilde{Z}(t, \beta) \geq \exp(-c(\log t)^5 t^{1/3}). \tag{3.13}$$

Now, we claim that

$$\mathbb{P}(R_t = G, \mathcal{E}_t) \leq \exp(-\frac{1}{4}\beta|\partial G|). \tag{3.14}$$

For each $x \in \partial G$, let y be a neighbour of x such that $y \notin G$. Consider the event $J_{xy}(t)$ that by time t there has never been a jump from x to y. On $\mathcal{E}_t$, x is visited at least β units of time. While at x, the rate of jumping to y is of course $1/4$. Let E_{xy} be independent exponential random variables with rate $1/4$, which represents the amount of time a particle would have to wait before jumping to y. Thus $J_{xy}(t) \cap \mathcal{E}_t \subset \{E_{xy} > \beta\}$. Hence

$$\mathbb{P}(R_t = G, \mathcal{E}_t) \leq \mathbb{P}(\cap_{x \in \partial G} J_{xy}(t) \cap \mathcal{E}_t) \leq \mathbb{P}(\cap_{x \in \partial G} E_{xy} > \beta) \leq e^{-\beta|\partial G|/4}$$

by independence of the random variables E_{xy}. Thus (3.14) is established.

Putting together (3.12) and (3.14) (resp. (3.13) and the definition of $\tilde{\mu}$), the proof of Theorem 1.4 (resp. Theorem 1.3) proceeds exactly as in Theorem 3.6.

4 Open problems and conjectures

We now discuss several open problems. The most basic ones are the following.

Problem 1. *Does Theorem 1.1 stay true if* $0 < \beta \leq \beta_0$*?*

It is possible to imagine that there is a phase transition from subdiffusive to diffusive as β decreases to 0. However simulations suggest that no such phase transition occurs and that the result should remain true even if $\beta \leq \beta_0$.

Our next question is to ask whether logarithmic factors in Theorem 1.1 are really present.

Problem 2. *Find* $f(t)$ *such that* $\operatorname{diam}(R_t)/f(t)$ *converges to 1 in* μ*-probability as* $t \to \infty$.

Another immediate question is to obtain the limiting shape of the random walk:

Problem 3. *Find a set* $S = S(\beta) \subset \mathbb{R}^2$ *such that*

$$\inf_{z \in \mathbb{R}^2} d_{Haus}\Big(\frac{R_t}{\operatorname{diam}(R_t)}; z + S\Big) \to 0$$

in probability, where d_{Haus} *stands for Hausdorff distance.*

The reason for looking for sets of the form $z + S$ to approximate the normalised range of the walk, is that we believe that with this formulation S can be chosen to be nonrandom: that is, a certain random translation of the deterministic shape S gives a good approximation of the normalised range.

Once the existence of S is established one may ask numerous questions about its geometry. For instance, does it have any flat facet? We conjecture in fact that as $\beta \to \infty$, $S(\beta)$ converges in the Hausdorff sense to a diamond of unit diameter. Indeed it is easy to see that the diamond is the minimiser of the isoperimetric problem for the vertex-boundary: i.e., if $\partial_v S = \{x \in S : \exists y \notin S, x \sim y\}$, then

$$\min_{|S| = v} |\partial_v S|$$

is attained for a diamond $S = \{x, y : |x| + |y| \le n\}$, whenever $v = 2n(n+1)$.

It is interesting to compare this situation for the case of a random walk conditioned on $\{L_t(x) \ge \beta, \forall x \in R_t\}$. There again, we believe a similar shape theorem holds, but the limiting shape of $S(\beta)$ as $\beta \to \infty$ should instead be a square with unit diameter. This is because, (3.14) in fact shows that

$$\mathbb{P}[R_t = G, \mathcal{E}_t] \le \exp(-\beta |\partial_e G|)$$

where $\partial_e G$ denotes the edge boundary of a graph G. Thus, when $\beta \to \infty$, it is reasonable to guess that $S(\beta)$ should minimise its edge boundary, rather than its vertex boundary , and hence be a square rather than a diamond.

Typical fluctuations of the shape (in the case where $\beta > \beta_0$ say) raise intriguing questions. In the percolation construction of the Wulff crystal when β is large but finite, these fluctuations are known with considerable precision. For instance, the so-called maximal local roughness, which measures the maximal distance from a point on the boundary of the shape to the polygonal hull of that shape, is of order $n^{1/3}$ if the diameter of the Wulff shape is of order n: see [23], [1]; more recently an extremely precise result in this direction was recently established by Hammond [18]. These exponents are identical to those arising in the Kardar–Parisi–Zhang (KPZ) universality class. While the Wulff crystal is not believed to belong to this universality class, there is nevertheless an analogy between the two situations, a fact which guides the intuition in the approach of [18].

Problem 4. *What are the maximum local roughness and facet length of the shape* R_t *when* $t \to \infty$*?*

We speculate that the exponent for the fluctuations take the value in agreement with the Wulff crystal (and hence also with KPZ): this suggests that the maximum facet length is of order $t^{2/9}$ and the maximum local roughness of order $t^{1/9}$.

Problem 5. *What is the diameter* $\operatorname{diam}(R_t)$ *when* $d \geq 3$*?*

We believe that, at least for β sufficiently large, the diameter is essentially given by the following formula:

$$\operatorname{diam}(R_t) \approx (t/\beta)^{\frac{1}{d+1}} \tag{4.1}$$

with high probability, possibly up to logarithmic factors. To see where this comes from, note that in d dimensions, the probability that a random walk stays in a box of side length L is approximately $\exp(-ct/L^2)$. The value of the energy should then be not too different from βL^{d-1}. Equating

$$\frac{t}{L^2} = \beta L^{d-1}$$

yields the value in the right hand side of (4.1).

Naturally, what makes this problem more difficult in three dimensions than two is that the topology of the boundary is much more complicated. For instance, the shape could conceivably have a few thin protuberances ("hairs") sticking out of a globally compact shape. Though each such occurrence increases the energy significantly and so should be penalised, the size of the ambient space available means that there are a lot of opportunities to do so. Such difficulty was already present in the study of the Wulff crystal for the Ising or percolation models (see e.g. [2]).

References

[1] K. Alexander. Cube-root boundary fluctuations for droplets in random cluster models. *Comm. Math. Phys.*, 224(3):733–781, 2001.

[2] K. Alexander, J.T. Chayes and L. Chayes. The Wulff construction and asymptotics of the finite cluster distribution for two-dimensional Bernoulli percolation. *Comm. Math. Phys.* 131, 1-50 (1990).

[3] E. Bolthausen. Localization of a two-dimensional random walk with an attractive path interaction. *Ann. Probab.*, 22, 875–918 (1994).

[4] R. Cerf. The Wulff crystal in Ising and Percolation models. *Lecture Notes in Mathematics* 1878, Springer. (Ecole d'été de Probabilités de Saint-Flour 2004.)

[5] P. Diaconis and L. Saloff-Coste. Comparison techniques for random walks on finite groups. *Ann. Probab.*, 21, 2131–2156 (1993).

[6] A. Dembo, Y. Peres, J. Rosen and O. Zeitouni. Thick points for planar Brownian motion and the Erdős-Taylor conjecture on random walk. *Acta Math.* 186 (2) 2001, 239–270.

[7] A. Dembo and O. Zeitouni. *Large Deviations Techniques and Applications.* Springer, Applications of Mathematics, Stochastic Modelling and Applied Probability, Second Edition.

[8] R. L. Dobrushin, R. Kotecký, S. B. Shlosman. Wulff crystal: a global shape from local interaction. AMS, Translations Of Mathematical Monographs 104, Providence, Rhode Island, 1992

[9] Donsker, M. D. and Varadhan, S. R. S. (1975). Asymptotic evaluation of certain Wiener integrals for large time. Proceedings of International Conference of Function Space Integration, Oxford, (1974), 15–33.

[10] Donsker, M. D. and Varadhan, S. R. S. (1975). Asymptotic evaluation of certain Markov process expectations for large time. I. *Comm. Pure Appl. Math.* 28 1–47.

[11] Donsker, M. D. and Varadhan, S. R. S. (1975). Asymptotic evaluation of certain Markov process expectations for large time. II. *Comm. Pure Appl. Math.* 28 279–301.

[12] Donsker, M. D. and Varadhan, S. R. S. (1976). Asymptotic evaluation of certain Markov process expectations for large time. III. *Comm. Pure Appl. Math.* 29 389–461.

[13] Donsker, M. D. and Varadhan, S. R. S., (1979). On the number of distinct points visited by a random walk. *Comm. Pure Appl. Math.* 32 721–747.

[14] Donsker, M. D. and Varadhan, S. R. S. (1983). Asymptotic evaluation of certain Markov process expectations for large time. IV. *Comm. Pure Appl. Math.* 36 183–212.

[15] P. Doyle and J.L. Snell. *Random walks and electric networks.* Available at arXiv:math.PR/0001057.

[16] Durrett, R. *Probability: Theory and Examples.* Duxbury Press, 1996.

[17] Grimmett, G. *Percolation.* Grundlehren der mathematischen Wissenschaften, vol 321, Springer, 1999 (second edition).

[18] A. Hammond. Phase separation in random cluster models I: uniform upper bounds on local deviation. *Comm. Math. Phys.*, to appear.

[19] Levin, A., Peres, Y. and Wilmer, E. *Markov chains and mixing times.* AMS lecture notes [online version].

[20] R. Lyons and Y. Peres. *Probability on trees and networks.* Book available on R. Lyons' website.

[21] Révész, P. *Random Walk in Random and Non-Random Environments.* World Scientific Publishing Co., 2005.

[22] Saloff-Coste, L. *Lectures on finite Markov chains.* Lectures on probability theory and statistics (Saint-Flour, 1996), Lecture Notes in Math., vol. 1665, Springer, Berlin, 1997, pp. 301–413. MR1490046 (99b:60119).

[23] H. Uzun and K. Alexander. Lower bounds for boundary roughness for droplets in Bernoulli percolation. *Probab. Theory Related Fields*, 127(1):62–88, 2003.

[24] Wulff, G. Zur Frage der Geschwindigkeit des Wachstums und der Auflösung der Kristallflächen. *Zeitschrift fur Krystallographie und Mineralogie*, 34 (1901), 5/6, 449–530.